



100

3

3

SLAC

Staff Scientist

MIRC

WPI-MANA

MANA

1cm²

100

0.1

100

40%

SLAC

1

WPI-MANA

0.1

100

1

100

2 40%

Communications Physics 2020 1 23
JSPS

[]

Communications Physics

1 23

Sub-molecular structural relaxation at a physisorbed interface with monolayer organic single-crystal semiconductors

Akifumi Yamamura, Hiromasa Fujii, Hirohito Ogasawara, Dennis Nordlund, Osamu Takahashi, Yutaro Kishi, Hiroyuki Ishii, Nobuhiko Kobayashi, Naoyuki Niitsu, Balthasar Blülle, Toshihiro Okamoto, Yusuke Wakabayashi*, Shun Watanabe*, and Jun Takeya*

DOI 10.1038/s42005-020-0285-7

1 23

7

1 23

10

TEL 04-7136-3790 / E-MAIL takeya@k.u-tokyo.ac.jp

TEL 04-7136-3788 / E-MAIL swatanabe@edu.k.u-tokyo.ac.jp

TEL 022-795-7750 / E-MAIL wakabayashi@tohoku.ac.jp

TEL 04-7136-5450 / E-MAIL info@edu.k.u-tokyo.ac.jp

TEL 022-795-6708 / E-MAIL sci-pr@mail.sci.tohoku.ac.jp

TEL 029-862-6216 / E-MAIL press-ml@aist.go.jp

TEL 082-424-3701 / E-MAIL koho@office.hiroshima-u.ac.jp

TEL 029-854-2040 / E-MAIL kohositu@un.tsukuba.ac.jp

TEL 029-879-6047 / E-MAIL press@kek.jp

(1)

28

6 1



(2)

IoT

10cm²/Vs

(3)

KEK


KEK

PF 2.5 GeV


PF-AR, 6.5 GeV

2

KEK




1



2 X

1

1



3 C₈-DNBDT-NW