

広島大学 インキュベーション研究拠点 The Research Center for Animal Science

日本型(発)畜産・酪農技術用発センター

特別セミナー

日時:2016年11月21日(月) 16:00~17:00

会場:広島大学 生物生產学部 C201教室

Heng-Yu Fan (Ph,D Professor)

(Life Science's Institute, Zhejiang University, Hangzhou, China)


Role of Newly Discovered Oocyte Factors in Regulating Maternal-zygotic Transition in Mammals.

The mRNAs stored in oocytes undergo general decay during maternal-zygotic transition (MZT), and their stability is tightly interconnected with meiotic cell cycle progression. We identified B-cell translocation gene-4 (BTG4) as an MZT licensing factor in mouse, BTG4 bridged CNOT7, a catalytic subunit of CCR4 NOT deadenylase, to eIF4E, a key translation initiation factor, and played a permissive role in maternal mRNA decay, Oocyte intrinsic MAPK cascade triggers translation of Btg4 mRNA stored in fully-grown oocytes by targeting its 3'-untranslated region, thereby couples CCR4 NOT deadenylase-mediated maternal mRNA decay with oocyte maturation and fertilization. On the other hand, maternally accumulated YAP in oocyte is crucial for zygotic genome

activation. These observations provide insights into the mechanisms of zygotic genome activation, and suggest potential implications of YAP activators in improving the developmental competence of cultured embryos in human assisted reproduction and animal biotechnology,

本セミナーは、

5研究科共同セミナーで



Structural & Molecular Biology 23, 387 394 (2016)