

---

---

## Signaling specificity of G protein-coupled receptor (GPCR): Lessons from melanin-concentrating hormone receptor 1

The signaling and trafficking of most GPCRs involves receptor interactions with G proteins, GRKs, arrestins and other receptors. In addition to these widespread GPCR associations, there are many other types of GPCR-interacting proteins that can interact with particular receptors to fine-tune receptor activity. Melanin-concentrating hormone (MCH) is a cyclic neuropeptide exerting its action through two GPCRs, MCHR1 and MCHR2. The extensive progress using genetic and pharmacological approaches has confirmed that the MCH-MCHR1 system is involved in feeding, energy homeostasis, sleep and emotional processing. In mammalian cells transfected with MCHR1, MCH is able to activate multiple signaling pathways including calcium mobilization, p-ERK activation, and inhibition of cyclic AMP generation through Gi/o- and Gq-coupled pathways. In this seminar, I review the current knowledge regarding: i) the common and distinguishing structural features of MCHR1 for receptor activation and G protein selectivity and ii) the mechanism controlling the fine-tuning of MCHR1 signaling by receptor-selective partners and their possible physiological significance. Lastly, we will discuss our very recent works on functional and physiological roles of MCHR1 selectively located in neuronal primary cilia.

### Reference:

#### Basic findings

Saito Y et al. *Nature* 400, 265, 1999

Saito Y et al. *J. Comp. Neurology*. 435, 26, 2001

#### Structure-Function relationship, GPCR fine-tuning

Tetsuka M et al. *Endocrinology* 145, 3712, 2004

Saito Y et al. *Endocrinology* 146, 3452, 2005

Miyamoto-Matsubara M et al. *Cellular Signaling*, 20, 2084, 2008

Miyamoto-Matsubara M et al. *Ann NY Acad Science*. 1200, 112, 2010

Hamamoto A et al. *Cellular Signaling*, 27, 818, 2015

#### GPCR localized in primary cilia

Hamamoto A et al. *Cellular Signaling*, 28, 572-584, 2016

Kobayashi et al. in press